

## CHAPTER 7

# Antibiotic Resistance in the Environment, with Particular Reference to MRSA

**William Gaze,\*<sup>1</sup> Colette O'Neill,\*  
Elizabeth Wellington,\* and Peter Hawkey<sup>†</sup>**

---

<b>Contents</b>		
	I. Introduction	250
	II. Evolution of Resistance	250
	A. Origins of antibiotic resistance genes	251
	B. Mechanisms of resistance	253
	III. Mechanisms of Horizontal Gene Transfer	254
	A. The role of integrons in resistance gene mobility	255
	B. Coselection for resistance genes	257
	IV. Antibiotics and Resistance Genes in the Environment	258
	A. Sewage sludge	258
	B. Farm animals	260
	C. Transfer from the environment to the clinic	262
	V. MRSA in the Nonclinical Environment	264
	A. Methicillin resistance in <i>Staphylococcus aureus</i>	264
	B. Environmental reservoirs of MRSA	265
	C. Pig associated MRSA	266
	D. Cattle associated MRSA	267
	E. Horse associated MRSA	268
	F. MRSA in companion animals	269
	VI. Conclusions	270
	References	270

\* Department of Biological Sciences, University of Warwick, Coventry CV47AL, United Kingdom

<sup>†</sup> Division of Immunity and Infection, The Medical School, Edgbaston, Birmingham B152TT, United Kingdom

<sup>1</sup> Corresponding author: Department of Biological Sciences, University of Warwick, Coventry CV47AL, United Kingdom

## I. INTRODUCTION

The introduction of  $\beta$ -lactam antibiotics (penicillins and cephalosporins) in the 1940s and 1950s probably represents the most important event in the battle against infection in human medicine. Even before widespread global use of penicillin, resistance was already recorded. *E. coli* producing a penicillinase was reported in Nature Journal in 1940 (Abraham and chain, 1940) and soon after a similar penicillinase was discovered in *Staphylococcus aureus* (Kirby, 1944). The appearance of these genes, so quickly after the discovery and before the widespread introduction of penicillin, clearly shows that the resistance genes pre-dated the clinical use of the antibiotic itself.


Intuitive reasoning would suggest that antibiotic resistance occurs because of direct selection produced by the use of antibiotics in humans and animals. For example, the mutations associated with increased resistance to fluoroquinolones have been documented in specific regions of the *gyrA*, *gyrB*, *griA*, and *griB* genes, which are referred to as the quinolone resistance-determining regions (QRDRs) (Piddock, 1998). Selection for resistance to a given antibiotic may take place within an infected human treated with antibiotics. However, selection may occur in other environments such as waste water treatment systems, agricultural environments where antibiotics may be of veterinary origin, or within an environmental background where antibiotic selection is provided by bacterial antibiotic producers.

In contrast to the scenario where resistance is conferred by mutation and selection by medical antibiotics, resistance can occur in an organism by the acquisition of a novel gene. New genes are acquired by horizontal gene transfer (HGT), through conjugation, transformation, or transduction. The origins of mobile antibiotic resistance genes may be from bacteria that have been subject to antibiotic selection in a nosocomial environment, or from environmental bacteria.

An example of an environment where HGT is likely to occur is soil. Practices such as sewage sludge and animal slurry application introduce complex mixtures of bacteria containing drug resistance genes, medical and veterinary antibiotics, and other chemicals such as detergents and surfactants to land, where interactions may occur with indigenous soil bacteria (Fig. 7.1).

## II. EVOLUTION OF RESISTANCE

Antibiotic resistance has two components: the evolution of genes with novel activities and the evolution of mechanisms allowing horizontal transfer throughout the microbial population.



**FIGURE 7.1** Anthropogenic sources of bacterial pathogens, pharmaceuticals, and heavy metals, which, in conjunction with indigenous soil bacteria, provide a mixture of genes and selective pressure for selection or coselection of antibiotic resistance.

### A. Origins of antibiotic resistance genes

Many modern  $\beta$ -lactam antibiotics such as the 7- $\alpha$ -methoxycephalosporins are secondary metabolites of *Streptomyces* species; the majority of streptomycetes and other actinobacteria produce constitutively expressed  $\beta$ -lactamases, which have a high GC content ( $\geq 70\%$ ). However, the  $\beta$ -lactamases encountered in human and animal pathogens all have GC contents in the order of 45–65%, suggesting a Gram negative origin. There are several Gram negative bacteria that exist in close proximity to antibiotic producers in the rhizosphere, which calls for protection against the toxic metabolites of their neighbors. Certain plant pathogens and rhizobacteria such as *Erwinia*, *Serratia*, *Flavobacterium*, *Pseudomonas*, *Chromobacterium* and *Agrobacterium* sp. produce carbapenems,  $\beta$ -lactams, and monocyclic  $\beta$ -lactams (Jensen and Demain, 1995).


The rhizosphere is the soil compartment influenced by plant root metabolism, and is high in nutrients derived from root exudates consisting of compounds such as organic acids, sugars, amino acids, vitamins, and carbohydrates. The rhizosphere is an important niche for microorganisms involved in nutrient recycling and plant health. The resistance genes in the rhizosphere are likely to result from competition between microorganisms for colonization sites. Key mechanisms responsible for the selection of medically significant bacteria in the rhizosphere are discussed

in a recent review (Berg *et al.*, 2005). Many mechanisms involved in the interaction between antagonistic plant-associated bacteria and their host plants are similar to those responsible for bacterial pathogenicity, including pathogenicity in humans (Rahme *et al.*, 1995).

It has long been suspected that the environment constitutes a reservoir of novel antibiotic resistance genes, although its significance has been overlooked in favor of evolution of resistance within the clinical environment. Arguably one of the most clinically important groups of  $\beta$ -lactamases in Gram negative bacteria at the moment are the CTX-M family (Livermore and Hawkey, 2005). The identification of environmental progenitors of the extended-spectrum  $\beta$ -lactamase (ESBL) CTX-M enzymes, responsible for resistance to 3rd generation cephalosporins (3GCs), in bacteria of the genus *Kluyvera* clearly indicates the significance of the environment in the evolution of emerging antibiotic resistance determinants (Bonnet, 2004; Rodriguez *et al.*, 2004). *Kluyvera* sp. are rare human pathogens causing infections similar to *E. coli* and are more often found associated with plants. *K. ascorbata* has been shown to enhance plant growth, particularly in heavy metal contaminated soils (Burd *et al.*, 1998, 2000). *K. georgiana* produces KLUG-1, whose nucleic acid sequence clusters with the CTX-M-8 group (Fig. 7. 2). Sequence similarity between the genes suggests that the natural  $\beta$ -lactamases of *K. ascorbata* and *K. georgiana* are the progenitors of the CTX-M-2 and CTX-M-8 groups, respectively (Bonnet, 2004). Evidence suggests that the process of gene transfer from the chromosome of *Kluyvera* to other clinically important bacteria has occurred several times involving different mobile elements, such as the IS-10-like element found upstream of both KLUG-1 and CTX-M-8, and ISEcp1 found upstream of KLUA-1 and members of the CTX-M-2 group (Poirel *et al.*, 2001). ESBLs confer low level resistance to  $\beta$ -lactams, *K. cryocrescens* possessing the KLUC-1 ESBL conferred resistance only to cefotaxime, ceftriaxone, ceftiofime, and aztreonam when cloned into *E. coli* (Decousser *et al.*, 2001). It is probable that KLUC-1 is only weakly expressed in *K. cryocrescens*, but mutations in the promoter region would confer ESBL resistance. Biochemical analysis of KLUC-1 revealed that substrate specificity and substrate profile are similar to those reported for CTX-M enzymes.

ESBLs confer resistance to 3GCs, which are semisynthetic molecules that do not exist in nature as far as is known; however, their structure is basically that of a cephalosporin of which a number of naturally produced compounds exist in the environment. Very little is understood about the prevalence, ecological function, and diversity of ESBL genes in the horizontal gene pool in soil, whether they are in transiently resident human/animal derived bacteria, or in the chromosomes of bacteria permanently residing in the rhizosphere.

A fluoroquinolone resistance gene *qnrA* has recently been established as originating in the water-borne species *Shewanella algae*; this genetically mobile plasmid-borne gene has now moved into clinically important bacteria, which



**FIGURE 7.2** Unrooted tree illustrating the phylogenetic relationships of KLUC-1, KLUG-1, and KLUA-1, *Kluyvera* sp. chromosomal genes, which are the putative progenitors of the CTX-M-1, CTX-M-8, and CTX-M-2 groups of ESBL enzymes (neighbor joining tree, Trex).

again shows the importance of the natural environment as a reservoir of clinically important antibiotic resistance genes (Nordmann and Poirel, 2005). The *qnrA* genes are embedded in complex *sul1*-type integrons, which also carry CTX-M-2 and CTX-M-9 ESBLs.

## B. Mechanisms of resistance

There are four main types of antibiotic resistance in bacteria (Hawkey, 1998). Antibiotic modification allows retention of the same target as sensitive strains, but the antibiotic is modified before it reaches the target.

$\beta$ -lactamases enzymatically cleave the  $\beta$ -lactam ring, inactivating the antibiotic. Some bacteria either stop the antibiotic from entering into the cell or pump the compound out of the cell by efflux. Carbapenem  $\beta$ -lactam antibiotics enter Gram negative bacteria via a membrane protein known as porin, but resistant bacteria lack the specific D2 porin responsible for transport and are therefore resistant (Pirnay *et al.*, 2002). Efflux via a membrane pump is a common mechanism found in Gram negative and Gram positive bacteria, in which five different superfamilies of efflux pumps conferring antibiotic resistance have been reported (Mahamoud *et al.*, 2007).

Changes in the target site also produce resistance; the antibiotics are able to reach the target but are not able to inhibit the target because of changes in the molecule. Enterococci are regarded as inherently resistant to cephalosporins because the enzymes responsible for cell wall synthesis (peptidoglycan production), known as penicillin binding proteins (PBPs), have a low affinity for cephalosporins and are not inhibited (Hawkey, 1998). Resistance to  $\beta$ -lactam antibiotics in pneumococci is entirely due to the development of altered forms of the high-molecular-weight PBPs, which have decreased affinity for the antibiotics (Coffey *et al.*, 1995). Altered PBPs have emerged by recombinational events between the *pbp* genes of pneumococci and their homologs in closely related streptococcal species.

The fourth antibiotic resistance mechanism (usually an enzyme) is the production of an alternative target that is resistant to the antibiotic, whilst continuing production of the sensitive target. Methicillin resistant *Staphylococcus aureus* (MRSA) produces an alternative penicillin binding protein (PBP2a), which is encoded by *mecA* carried on the Staphylococcal Cassette Chromosome *mec* (SCC*mec*). Because PBP2a is not inhibited by the antibiotics, the cell continues to produce peptidoglycan and maintains a stable cell wall (Hardy *et al.*, 2004).

### III. MECHANISMS OF HORIZONTAL GENE TRANSFER

Gene transfer in the environment is central to the hypothesis that a reservoir of novel resistance genes exists outside the clinic, which can be transferred to clinically significant bacteria in hospitals. An extensive literature on genetic exchange between bacteria in the environment exists, which is reviewed elsewhere (Davidson, 1999). The current review concentrates specifically on gene transfer mediated by transposable elements such as class 1 integrons, which are of increasing clinical importance. The occurrence of coselection by nonantibiotic xenobiotics is also discussed.

The evolutionary response of bacteria exposed to antibiotics has been mediated in large part by the movement of conjugative plasmids carrying

antimicrobial resistance genes in both Gram negative and Gram positive bacteria. Transfer frequencies for conjugative plasmids in *Enterococcus faecium* can be as low as  $\geq 10^{-6}$  per donor in laboratory experiments using broth mating, but the production of short peptides, by some recipient cells, that cause cell aggregation greatly increases transfer rates ( $\geq 10^{-4}$ ) (Dunny *et al.*, 2001). Plasmids play an important role in gene transfer in staphylococci; conjugative plasmids are also capable of mobilizing small nontransferable plasmids (McDonnell *et al.*, 1983). Gene transfer does not always require the presence of plasmids, as the extensively studied and widely distributed conjugative transposons such as Tn916 demonstrate (Burrus and Waldor, 2004). The gut of both humans and food animals is an important site for transfer of conjugative plasmids, as careful studies have shown there is no loss of fertility of conjugative plasmids in the gut. Conjugation probably occurs continuously in the gut of humans/animals even in those not receiving antibiotics (Freter *et al.*, 1983). The release of faeces into the environment represents one of the most important sources of novel assortments and types of resistance one of genes in new bacterial hosts. The laboratory simulation of plasmid transfer in the environment can give low transfer frequencies, but these must be compared to the size of the populations involved. The requirement for close contact in the laboratory (e.g., filter mating) may readily be met in microenvironments: staphylococci on the skin, oral streptococci in dental plaque, or pseudomonads in water films on soil particles.

### A. The role of integrons in resistance gene mobility

In addition to multiresistance plasmids, antibiotic resistance genes are situated on transposable elements that can associate with other elements such as chromosomes. These transposable elements include transposons and integrons, which can be transferred horizontally. Integrons are recombination and expression systems that capture genes as part of a genetic element known as a gene cassette (Recchia and Hall, 1995). Gene cassettes bear a recombination site known as a 59-base element (59-be) that is recognized by the integron-encoded integrase IntI (Hall *et al.*, 1991). Most cassette genes described are antibiotic or quaternary ammonium compound resistance genes. However, recent studies have revealed that the cassette gene pool is far more diverse than previously thought. Stokes *et al.* (2001) designed PCR primers to conserved regions within the 59-be of gene cassettes, allowing detection of a large number of novel genes. Using these primers, 123 cassette types were recovered from Antarctic and Australian soils and sediments, with very few represented in clone libraries more than once indicating the large size of the cassette gene pool.

Most ORFs did not match known sequences, again illustrating the diversity of these gene sequences. Further studies revealed an additional 41 environmental gene cassettes, giving a total of 164 directly sampled from natural environments by PCR (Holmes *et al.*, 2003). There are several classes of integron, the most commonly studied being class 1 integrons that are commonly associated with antibiotic resistant bacteria. Recent studies have detected novel integron classes in soils, Nield *et al.* (2001) classified three new integron classes and Nemergut *et al.* (2004) an additional 14 classes, demonstrating the immense variety of these elements present in the environment.

The variable regions of class 1 integrons contain the cassette genes, and to the right of this lies the 3'-conserved region, which may have one of three different backbone structures (Partridge *et al.*, 2002). The first backbone type consists of a Tn402 (In16) like arrangement consisting of a *tni* module containing 3 transposition genes and a resolvase gene, the second In5 type consists of *qacEΔ1*, *sul1*, *orf5*, *orf6*, and a partial *tni* module (*tniΔ*) consisting of two transposition genes. The third In4 type just carries *qacEΔ1*, *sul1*, *orf5*, and *orf6*. Integrons carrying the complete *tni* module are able to undergo self-transposition and it is thought that In5 and In4 types may also be able to move if the *tni* gene products are supplied in *trans* (Partridge *et al.*, 2002).

The role of class 1 integrons in conferring antibiotic resistance to clinical isolates of many bacterial strains is well documented (Briggs and Fratamico, 1999; Leverstein-van Hall *et al.*, 2003; Segal *et al.*, 2003; White *et al.*, 2000). Fluit and Schmitz (2004) summarized recently described cassette gene diversity which includes 25 β-lactam resistance genes (including 8 carbapenemase and 17 ESBL genes), 11 aminoglycoside, 2 chloramphenicol, rifampicin, 3 trimethoprim, and quinolone resistance genes. It is therefore clear that integrons are capable of conferring resistance to extended-spectrum β-lactams, carbapenems and fluoroquinolones, representing an extremely efficient method of acquiring resistance to the most widely used and important antibiotics. Studies on the incidence of class 1 integrons in bacterial pathogens associated with agriculture and fish farming such as *E. coli* and *Aeromonas salmonicida* have shown a link between integrons and antibiotic resistance (Bass *et al.*, 1999; Sorum *et al.*, 2003). A recent study (Nemergut *et al.*, 2004) illustrated evidence of a gene cassette encoding nitroaromatic catabolism, a group of compounds associated with mining activity, which highlights the fact that selective pressures other than antibiotics may also coselect for resistance genes. The process by which a gene becomes a movable cassette is not understood; however, it has been proposed that the 59-be is added as a transcription terminator to an RNA gene transcript, which is subsequently converted into DNA by a hypothetical reverse transcriptase (Recchia and Hall, 1995).



Class 1 and class 2 (also capable of carrying antibiotic resistance genes) are known to undergo transfer between bacteria in chicken litter, which is often spread onto soil where further selection and horizontal transfer may occur (Nandi *et al.*, 2004). The integrons study was the first to demonstrate wide spread prevalence of class 1 integrons in Gram positive bacteria, illustrating the potential for HGT.

## B. Coselection for resistance genes

It is commonly assumed that in the absence of antibiotic selection mobile resistance genes will be lost, and the host return to a sensitive phenotype, as the genes confer a metabolic cost on the host. Coselection is one mechanism whereby other resistance genes carried on the same genetic element produce selection for an entire mobile genetic element. Anthropogenic activity produces emissions of complex mixtures of xenobiotics, bacterial pathogens, and antibiotic resistance genes into the environment in the form of industrial and domestic effluent as well as human and animal waste. Industrial and domestic pollutants such as quaternary ammonium compounds (QACs) have been shown to exert an extremely strong selective pressure for class 1 integrons, which are a major mechanism for dissemination of antibiotic resistance (Gaze *et al.*, 2005). Coselection is produced by the presence of QAC resistance genes on multiresistance plasmids or class 1 integrons. QAC resistance genes fall into two families: *qacA/B* belong to the Major Facilitator Superfamily and are only found in staphylococci on multiresistance plasmids (Paulsen *et al.*, 1996). Other QAC resistance genes belong to the Small Multidrug Resistance Family and include *qacC/D* now known as *smr*, *qacE*, *qacEΔ1*, *qacF*, *qacG*, *qacH*, and *qacJ* (Gaze *et al.*, 2005). *QacE*, *qacEΔ1*, *qacF*, and *qacG* have been identified on integrons and the remaining genes on multiresistance plasmids in staphylococci. In a study investigating QAC pollution and class 1 integron prevalence, bacteria were isolated from a reed bed used to remediate effluent from a textile mill (Gaze *et al.*, 2005). QAC resistance was higher in isolates from reed bed samples and class 1 integron prevalence was significantly higher in populations pre-exposed to QACs.

Exogenous plasmid isolation can be used to detect resistance genes in soil bacteria. This method allows capture of plasmids from the total bacterial fraction of an environmental sample without the necessity to culture the host organism. Smit *et al.* (1998) investigated mercury resistance plasmids in soil populations using exogenous isolation, and identified plasmids of 10–50 kb carrying resistance to copper, streptomycin, and chloramphenicol. These authors amended soil with mercuric chloride and found this to subsequently increase the recovery of resistance plasmids, highlighting the fact that heavy metals may coselect for antibiotic resistance in the environment. Plasmids have also been captured from

polluted soils and slurries (Smalla *et al.*, 2000; Top *et al.*, 1994). The latter authors identified multiple antibiotic resistance genes from isolated plasmids. Some aspects of animal husbandry using heavy metal containing compounds select for antibiotic resistance genes in the environment, for example, the use of copper growth supplements in pigs (Hasman *et al.*, 2006). In Denmark glycopeptides were banned in food animal production in 1995, following that ban the rate of glycopeptide resistance in *Enterococcus faecium* (GRE) did not change. The banning of macrolides in 1998 led to a significant fall in glycopeptide resistance, this was thought to be due to both antibiotic resistance genes being on the same plasmid and coselection by the continuing use of macrolides (Aarestrup, 2000). GRE continue to be prevalent in Danish pigs; the reason for this is now thought to be the linkage on the same plasmid of the copper resistance gene *tcrB* and macrolide/glycopeptide resistances. A feeding experiment using 175 mg Cu/kg in feed selected for GRE in piglets, whereas 6 mg/kg feed did not (Hasman *et al.*, 2006).

#### IV. ANTIBIOTICS AND RESISTANCE GENES IN THE ENVIRONMENT

Human and animal wastes may contain antibiotics or active intermediates from human and veterinary medicines that may potentially increase antibiotic resistance selection in soil, in addition to introducing pathogens, which can exchange mobile genes with indigenous rhizosphere bacteria. Antibiotics retain their selective capabilities in the soil and are ultimately released to surface waters (Boxall *et al.*, 2002). Certain plant pathogens and rhizobacteria such as *Erwinia*, *Serratia*, *Flavobacterium*, *Pseudomonas*, *Chromobacterium*, and *Agrobacterium* sp. produce carbapenems,  $\beta$ -lactams, and monocyclic  $\beta$ -lactams (Jensen and Demain, 1995). In addition, streptomycetes and fungi produce a wide range of antibiotics.

##### A. Sewage sludge

The 2001 UK Sewage Sludge Survey showed that an average of 1,072,000 tonnes of dry solids per annum was produced in 1998–2000. The UK Department for Food and Rural Affairs (DEFRA) state that conventionally treated sludge has been subjected to defined treatment processes and standards that ensure at least 99% of pathogens have been destroyed. Enhanced treatment, originally referred to as “Advanced Treatment,” describes treatment processes capable of virtually eliminating any pathogens (99.9999%) that may be present in the original sludge. Conventionally treated sewage sludge cannot be surface spread if the land is intended for grazing; it must be deeply injected into the soil and

left for at least three weeks until it is grazed. Conventionally treated sewage can be applied to the surface of grassland, or for forage crops such as maize, which will then be harvested (no grazing allowed within the season of application). If applied to vegetable-growing land at least 12 months must have elapsed between treatment and harvest. When using enhanced treated sludges farmers must wait at least three weeks before grazing the animals or harvesting the forage crops, and at least 10 months before harvesting fruit and vegetable crops grown in direct contact with the soil and normally eaten raw (<http://www.defra.gov.uk/farm/waste/sludge/index.htm>). It is clear that the potential for transmission of resistance genes to food animals and vegetable crops exists.

A hydraulic, biokinetic, and thermodynamic model of pathogen inactivation during anaerobic digestion showed that a 2 log<sub>10</sub> reduction in *E. coli* (the minimum removal required by the UK government for agricultural use of conventionally treated biosolids) is likely to challenge most conventional mesophilic digesters (Smith *et al.*, 2005). UK regulations for pathogen removal are becoming more stringent, but the processes used to reduce bacterial indicator species numbers may have a quite different effect on resistance gene numbers.  $\beta$ -lactam and aminoglycoside resistance genes have been isolated by exogenous isolation from activated sewage in Germany, illustrating that final stage sludge is a source of antibiotic resistance genes (Tennstedt *et al.*, 2005).

Crucially, although sewage sludge has been demonstrated to contain antibiotic resistance genes and pathogenic bacteria, the extent of this problem and the potential for transfer of resistance to soil bacteria and ultimately its effect on the human population is unknown. Sewage sludge also contains measurable concentrations of antibiotics which may continue to select for resistance in the soil. A study by Golet *et al.* (2003) suggested that sewage sludge is the main reservoir of fluoroquinolone (FQ) residues from waste waters and outlined the importance of sludge management strategies to determine whether most of the human-excreted FQs enter the environment. Field experiments of sludge-application to agricultural land confirmed the long-term persistence of trace amounts of FQs in sludge-treated soils and indicated a limited mobility of FQs into the subsoil. Where sewage treatment plants receive large amounts of effluent from hospitals, the problem of antibiotic residues and resistance genes in sewage sludge may be particularly significant. Persistence of FQs is particularly relevant as they appear to coselect for class 1 integrons and integron borne ESBL genes. The recent discovery of new quinolone resistance genes (Nordmann and Poirel, 2005), which are situated on class 1 integron structures and also confer ESBL resistance, reinforces this. Our own research has indicated an impressive reservoir of FQ resistance (QRDR mutations) in staphylococci associated with free range chicken farming (Hawes, 2004).

Recent studies in Portugal have identified  $\beta$ -lactamases, including TEM, IMP, and OXA-2 derivatives in aquatic systems, and ESBL resistance genes in sewage sludge, which is spread to land in the UK and therefore has the potential to recycle resistance to the human food chain (Henriques *et al.*, 2006). ESBL-producing Enterobacteriaceae were detected in five samples of human sewage in Spain (Mesa *et al.*, 2006).

The human colon is the major reservoir of emerging opportunistic pathogens such as *E. coli*, *Klebsiella*, *Enterobacter*, and *Acinetobacter baumannii* (Agustía *et al.*, 2002; Fanaro *et al.*, 2003; Hollander *et al.*, 2001) and it is likely that these are food-derived in the community (Turtura *et al.*, 1990). The distinction between food-borne commensals, pathogens and nosocomial pathogens is somewhat arbitrary, and many emerging nosocomial Gram negative pathogens may be food borne, normally living in the gut as commensals until the individual becomes immunosuppressed or until antibiotic resistance genes transferred from another organism. Recent research has revealed that soil, particularly the plant rhizosphere, harbors diverse opportunistic human pathogens, including *Acinetobacter baumannii*, *Aeromonas salmonicida*, *Burkholderia cepacia*, *Enterobacter agglomerans*, *Klebsiella pneumonia*, *Proteus vulgaris*, *Pseudomonas aeruginosa*, *Salmonella typhimurium*, *Serratia sp.*, *Staphylococcus aureus*, *Stenotrophomonas maltophilia*, and others (Berg *et al.*, 2005).

## B. Farm animals

When soils are treated with manure both the residues of antibiotics used as veterinary medicines and the bacteria carrying genes conferring resistance are introduced into the soil and reach the food chain, for example, via plant-associated bacteria (Witte *et al.*, 2000). Some 461 tonnes (active ingredient) of antimicrobial therapeutics and growth promoters were sold for use in food animals in 2000 in the UK (National Office of Animal Health data), including tetracyclines (228 t), trimethoprim/sulphonamides (94 t),  $\beta$ -lactams (49 t), macrolides (41 t), and FQs (1 t). Some veterinary antibiotics are synthetic, so unlike those produced by soil bacteria, many cannot be broken down through normal processes, and therefore may persist for a long time in soils; adsorption to soil particles and other surfaces allows accumulation of residual antibiotics to high concentrations (Kummerer, 2004).

It has been clearly established that the use of certain antibiotics in agriculture has contributed to the development of resistant bacterial strains in human infection, as evidenced by the work of Wolfgang Witte and others where vancomycin resistance genes in human *Enterococcus faecalis* isolates were traced to the use of avoparcin in pigs (Witte, 1997). As early as the 1970s Stuart Levy noted that oxytetracycline was a major feed additive and that studies had shown there was a strong association between tetracycline resistance in isolates from livestock and animal

workers (Levy *et al.*, 1976). In 1969 the Swann committee report recommended that antibiotics used in human medicine should not be used as growth promoters. Only in the last few years have a number of antibiotics been banned, and clearly their use as growth promoters is inadvisable.

A mixture of the parent product and metabolites are excreted in faeces and urine. Excreta enter the farm environment directly in grazing animals and indirectly in intensively reared animals through application of slurry and manure. It is estimated that approximately 70 million tons of animal manure wastes are spread onto agricultural land per annum in the UK (Hutchison *et al.*, 2004). In many river catchments the bulk of faecal coliforms are believed to be of agricultural origin. Chee-Sanford *et al.* (2001) screened for eight *tet* genes in groundwater associated with swine production facilities. Tetracycline resistance genes were found as far as 250 m downstream from waste lagoons, highlighting the danger posed by use of antibiotics in agriculture and the risk of contamination of drinking water with antibiotic resistant bacteria. In a different study a detection limit of  $10^2$ – $10^3$  copies of the *tet*(M) gene per gram was achieved using a nested PCR method with TC-DNA (Agero *et al.*, 2004). The gene was detected in farmland soil previously amended with pig slurry containing resistant bacteria; the number of positive samples from farmland soils one year after manure treatment was significantly higher than in samples of garden soil not treated with manure.

*E. coli* strains producing an ESBL (CTX-M-2) were recently isolated from cattle faeces in Japan (Shiraki *et al.*, 2004).  $\beta$ -lactamases and ESBLs were also detected in *E. coli* isolates from healthy chickens, food and sick animals in Spain (Brinas *et al.*, 2002, 2003a,b). The use of extended-spectrum cephalosporins in chickens is very unusual, and the possibility of cross-selection with other antimicrobials used in poultry (such as sulphonamides and tetracyclines, etc.) might explain this discovery. In a study of retail chicken breasts, quinolone resistant *E. coli* also producing CTX-M-2 were found in 5 of 10 samples produced in Brazil (CTX-M-2 is widely reported in human infection with *Salmonella* and *E. coli* from South America), whereas only 1 of 62 samples of UK produced chicken were positive for CTX-M-1 (Ensor *et al.*, 2007). ESBLs were also discovered in samples from 8 of 10 pig farms, 2 of 10 rabbit farms, from all 10 poultry farms tested and in 3 of 738 food samples studied in Spain (Mesa *et al.*, 2006).

The FQ antibiotic enrofloxacin (the major metabolite being ciprofloxacin) is used extensively in poultry farming and evolution of fluoroquinolone resistance in chicken litter was documented, caused by mutation in the QRDR of the *gyrA* gene (Lee *et al.*, 2005). Plasmid-mediated quinolone resistance (PMQR) is also known to occur. The gene responsible for PMQR was identified as *qnr* and this gene was found in an integron-like element and is associated with the ESBL VEB-1 (Poirel *et al.*, 2005c). The origin of *qnrA* was recently established as the water-borne species

*Shewanella algae*, which, in addition to the fact that CTX-M-ases originate in the rhizosphere bacteria *Kluyvera* sp., underlines the role of the environment itself as a reservoir of novel resistance genes (Nordmann and Poirel, 2005; Poirel *et al.*, 2005b). Recently, further transferable quinolone resistance genes *qnrB* and *qnrS* have been identified in Gram negative opportunistic pathogens (Poirel *et al.*, 2005a). In 2004, an entirely novel plasmid mediated mechanism of quinolone resistance was discovered. In *E. coli* from Shanghai, PRC strains with a MIC of 1.0 mg/liter of ciprofloxacin carried a mutated form of the aminoglycoside inactivating enzyme AAC (6<sup>1</sup>)-Ib-cr (Robicsek *et al.*, 2006). This enzyme N-acetylates quinolones that have an amino nitrogen on the piperazinyl substituent (e.g., ciprofloxacin and norfloxacin). The distribution of the gene has not been studied extensively but has been found in China and North America (Robicsek *et al.*, 2006). It was recently reported to be associated with the ESBL genes *bla*<sub>CTX-M-15</sub>, *bla*<sub>OXA-1</sub>, and *bla*<sub>TEM-1</sub> on a limited range of plasmids carried by *E. coli* in Portugal (Machado *et al.*, 2006). The consequences could be grave as ciprofloxacin is extensively used in agriculture and is the principal metabolite of veterinary quinolones such as enrofloxacin. The use of quinolones (which are also slow to degrade in the environment) will coselect for a wide range of *qnr/acc* (6<sup>1</sup>) associated resistances such as resistance to 3rd generation cephalosporins and carbapenems. Thus, there is increasing evidence of an environmental reservoir of clinically important antibiotic resistance genes.

Several studies demonstrated a correlation between the extent of use of antibiotics in animals and the incidence of the respective antibiotic resistance genes. Therefore it is not surprising to learn that animal manure, in particular piggery manure, has been shown to be a "hot spot" (high incidence) of bacteria carrying antibiotic resistance genes residing on mobile genetic elements (Smalla *et al.*, 2000).

Antibiotic resistance does not always appear to be directly related to short-term trends in antibiotic usage. The glycopeptide growth promoter avoparcin was banned from animal production in Denmark in 1995, and in the EU in 1997, because of concern for the spread of vancomycin-resistant enterococci (VRE) from food animals to humans. A Danish study found high levels of VRE in broiler flocks five years after avoparcin was withdrawn (Heuer *et al.*, 2002a). Further studies revealed that VRE was surviving in broiler houses despite cleaning procedures between production rotations (Heuer *et al.*, 2002b).

### C. Transfer from the environment to the clinic

Patients admitted to hospital are likely to acquire bacteria that are multiplying in the hospital environment, such as *Pseudomonas aeruginosa* and these may well be antimicrobial resistant. It has been demonstrated that

the general environment of the hospital, particularly sites such as sink drains, mop heads, and other wet environments, will act as sources not only of bacteria capable of directly causing nosocomial infection but which can also act as a gene pool of antibiotic resistance genes. Apart from this obvious interaction of patients with the hospital environment, it is probably food that is the major route of flow of resistance genes from the more general environment to man. The human gut carries a large number of commensal bacteria, usually in the order of  $\geq 10^{10}$ /g of faeces and this value applies only to culturable bacteria. The bowel flora is constantly being challenged by new bacteria in food and although the dominant flora remains, colonization with a minority of antibiotic resistant bacteria, particularly Enterobacteriaceae/enterococci and staphylococci, occurs in those individuals not receiving antimicrobials. The size and complexity of this bowel flora reservoir, which is in direct connection with the environment, has long been recognized as containing large numbers of antibiotic resistant bacteria. A study undertaken as early as 1979 showed that in individuals with no history of recent consumption of antibiotics, 10% or more of the total aerobic Gram negative bacteria were resistant to one or more antimicrobials (Levy *et al.*, 1988). This pool of resistance genes may get transferred into bacteria with significant pathogenicity towards humans, or in very many cases, opportunistic pathogenic bacteria such as *E. coli*, *Klebsiella* sp., enterococci, and *Staphylococcus aureus*, which may cause infections in the individual. In this way, there is a continuous link between selection for antibiotic resistance in the general and agricultural environment and human medicine. The other major commensal flora site on humans is the skin surface and recently with the recognition of community acquired MRSA, the importance of antibiotic resistant staphylococci and other Gram positive bacteria on the skin, and the transfer into potentially pathogenic species such as *Staphylococcus aureus* has been emphasized. As with the large bowel, all the studies clearly show that even those individuals not receiving antibiotics carry a substantial load of antibiotic resistant bacteria, which must be strongly influenced by interactions with other humans, companion and food animals. An extensive study by Cove and colleagues showed that the incidence of seven primary antibiotic resistance markers among the staphylococcal flora in antibiotic untreated subjects was tetracycline 87.5%, erythromycin 68.8%, fusidic acid 56.3%, trimethoprim 42.4%, chloramphenicol 25%, clindamycin 9.4%, and gentamicin 4.7%. We now recognize that among staphylococci there is ample opportunity and genetic mechanisms for the mobilization of resistance genes into potentially pathogenic species (Cove *et al.*, 1990). Some special groups in the human community such as farm workers, medical personnel and patients receiving antimicrobials, have a much higher incidence of colonization with antibiotic resistant bacteria, some of which, such as in the case of farm workers, are

derived from contact with farm animals that have either been treated with antimicrobials or exposed to selecting agents. Recent problems with community acquired MRSA in pig farmers illustrate the way in which problems can arise rapidly, and undermine the use of clinically important antimicrobials.

## V. MRSA IN THE NONCLINICAL ENVIRONMENT

### A. Methicillin resistance in *Staphylococcus aureus*

*Staphylococcus aureus* is well known for its ability to acquire antibiotic resistance, both historically in relation to penicillin, erythromycin, and tetracycline and more recently methicillin and vancomycin resistance. The acronym MRSA (Methicillin resistant *S. aureus*) is feared by health-care professionals the world over. *S. aureus* forms part of the normal human flora, residing asymptotically in the mucosal linings of healthy individuals and at other moist skin sites (Hiramatsu *et al.*, 2001; Peacock *et al.*, 2001) and is particularly pathogenic in individuals at the extremes of age who have intravascular/urinary catheters, diabetes, and other compromising coexisting medical conditions (Lindsay and Holden, 2004). However, recent years have seen a rise in highly virulent community acquired strains (CA-MRSA) capable of causing disease in young, healthy individuals with none of the prescribed risk factors. Resistance to methicillin is carried by *SCCmec*, a mobile genetic island that can disseminate horizontally, although its mode of transfer is currently unknown (Hanssen and Ericson Sollid, 2006). Resistance is encoded by the *mecA* gene (Ito *et al.*, 1999), *mecA* encodes an attenuated penicillin binding protein (PBP2' or PBP2a), which has a lower affinity for penicillin and other  $\beta$ -lactams than the innate PBPs (Hartman and Tomasz, 1981), hence interfering with antimicrobial activity. There are six basic described *SCCmec* types; additional resistance genes may be present or absent depending on the type. *SCCmec* is inserted into the *S. aureus* chromosome near the origin of replication, always at the same location at the 3' end of the *orfX* gene (Kuroda *et al.*, 2001). Its origins are unknown, but as no methicillin-susceptible *S. aureus* (MSSA) homolog exists (Archer and Niemeyer, 1994) it has been suggested that it was transferred horizontally from a coagulase-negative staphylococcus (CoNS) such as *S. sciuri* (Couto *et al.*, 1996, 2003). The conjugative transposon containing *tetM* (tetracycline resistance) has been suggested to pass between *Clostridia* and staphylococci (Ito *et al.*, 2003), and empirical support for the movement of mobile elements between staphylococci and other low GC Gram negative bacteria (Gill *et al.*, 2005), and the potential transfer of *vanA* (vancomycin resistance) from *Enterococcus faecalis* to *S. aureus* (Weigel *et al.*, 2003)



lends weight to the possibility of a common gene pool available to many bacterial species (Hanssen *et al.*, 2004). Movement of DNA into *S. aureus* is tightly controlled by a restriction-modification system Sau1, a type 1 system that has been found on the chromosome of all known sequenced strains (Waldron and Lindsay, 2006).

Although much of the genome of *S. aureus* consists of mobile genetic elements, because of the presence of Sau1, horizontal transfer is likely to be more frequent among members of the same lineage, as the Sau1 restriction-modification system present in different lineages have specific differences (Waldron and Lindsay, 2006). This is thought to explain the rare occurrence of *vanA* carrying VRSA and the limited number of lineages with SCC*mec* (CC1, CC5, CC8, CC22, CC30 and CC45). Possession of SCC*mec* is thought to carry a fitness cost and will therefore be selected for only when strains have been exposed to antibiotics (Katayama *et al.*, 2003).

## B. Environmental reservoirs of MRSA

Although antibiotic selective pressure present in clinical environments is clearly a major source of MRSA infection (Dar *et al.*, 2006), environmental reservoirs have been implicated in the spread of resistant strains. Subinhibitory levels of antibiotics may be to blame for inducing resistance in commensal bacteria in farm animals, inducing resistance in pathogenic bacteria through plasmid transfer (Singer *et al.*, 2003). In pig and poultry farming, heavy use of antibiotics in typically intensively farmed settings predisposes them to MRSA colonization (Shea, 2004; van Den Bogaard *et al.*, 2000). Close contact between animals is inevitable, and infections are likely to spread quickly. Bacterial transmission in humans is (or should be) easily negated through regular hand washing, whereas oral-faecal contact cannot be prevented in animals and fast transmission of faecal-borne disease is unavoidable (van den Bogaard and Stobberingh, 1999). Therefore, an infection in one or two animals is often combated by a blanket treatment with antibiotics of the entire house, which may contain over 10,000 animals (Shea, 2004). The farming practice employed may have some bearing on the antibiotic susceptibility of commensal *S. aureus*; organic farming may result in fewer resistant bacteria than in conventional farms because of lower exposure to antibiotics and reduced contact between animals (Halbert *et al.*, 2006; Sato *et al.*, 2005; Tikofsky *et al.*, 2003). There is some contrary evidence (Sato *et al.*, 2004), but the continued use of antibiotics in some of the organic farms studied was a possibility (Busato *et al.*, 2000).

Cross-resistance for methicillin and some cephalosporins has been demonstrated (Hansen-Nord *et al.*, 1988; Menzies *et al.*, 1987), with administration of cephalosporins increasing the acquisition of nosocomial

MRSA three-fold (Asensio *et al.*, 1996). Similarly, tetracycline resistance may be caused by one of several genes, one of which, *tetK*, is carried by a plasmid (pT181) that is inserted into SCC*mec*III (Ito *et al.*, 2003). As tetracycline concentrations have been found to persist at high concentrations long after slurry application has ceased (De Liguoro *et al.*, 2003; Hamscher *et al.*, 2002), selection for tetracycline resistance may contribute to methicillin resistance as a result of selection for SCC*mec*III; although SCC*mec*IV is far more common in community MRSA. In the absence of antibiotics, the presence of quaternary ammonium compounds (QACs) in soils, used extensively as disinfectants, may also coselect for  $\beta$ -lactam resistance (Sidhu *et al.*, 2001, 2002); resistance to QACs correlates with the  $\beta$ -lactamase transposon Tn552 because of co-carriage of the *blaZ* gene and *qacA* (Anthonisen *et al.*, 2002). Evidently, the introduction of any of these compounds to the environment may lead to selection for methicillin resistance in *S. aureus* colonizing or infecting farm animals, or in other bacteria inhabiting the soil; there is some evidence to suggest colonization of the rhizosphere by *S. aureus* may be possible, so acquisition of methicillin resistance may occur both in soil and in farm animals (Berg *et al.*, 2005; Germida and Siciliano, 2001; Morales *et al.*, 1996).

### C. Pig associated MRSA

Antibiotic resistance levels vary from country to country. For example, the percentage of MRSA in the UK (40% of nosocomial *S. aureus* isolates causing bacteremia) contrasts sharply with that in Holland, where it is currently very rare, making up only 1% of *S. aureus* isolates (Tiemersma *et al.*, 2004). In Holland most cases are found in people who have recently attended foreign medical institutions; however, occasional instances have arisen, whereby foreign travel was not a contributing factor. One such case described an MRSA positive baby that had never travelled abroad (Voss *et al.*, 2005). The family lived on a pig farm, and MRSA was isolated from one of the pigs. Two further cases were reported in the same study, both of which were associated in some way with pig farming. As pig farming was the only common denominator in these cases it was deduced that there may be a link between pig farming and increased risk of MRSA infection; indeed colonization by MRSA in pig farmers was found to be almost twice that of the general population (Aubry-Damon *et al.*, 2004). Pigs have been similarly implicated as a reservoir of MRSA in France, a study of pig isolates found all to be nontypeable by pulsed-field gel electrophoresis (PFGE), the so-called gold standard for MRSA typing (Armand-Lefevre *et al.*, 2005), and the association between PFGE nontypeable strains and pig farming has gained further empirical support (Huijsdens *et al.*, 2006). The three multilocus sequence types (MLSTs) ST9, ST398, and ST433 were found in pigs and only in humans associated

with pig farming suggesting an association between these STs and pigs (Armand-Lefevre *et al.*, 2005). These findings were supported by a recent Dutch study in which 39% of all pigs sampled from slaughter houses (540 pigs) carried MRSA in their nares, and all of these belonged to ST398 (de Neeling *et al.*, 2007). Obviously Holland represents an unusual case because of its low MRSA incidence, but the high percentage of swine carriers, all with the same MLST type, suggests a recent dissemination of MRSA. The vast majority of isolates contained SCC*mec* types IV and V, indicating a relatively frequent transfer of this mobile element; types IV and V are suspected to transfer at a greater rate than the other types (Daum *et al.*, 2002). As pig-associated strains seem to be passed from human to human as well as between humans and pigs, pig farming may represent a source of CA-MRSA (Huijsdens *et al.*, 2006). CA-MRSA is susceptible to a greater range of antibiotics than its hospital-acquired counterpart (Herold *et al.*, 1998), but tends to be more virulent because of the acquisition of toxin genes such as the Panton-Valentine Leukocidin (PVL) (Boyle-Vavra and Daum, 2007). PVL has not yet been found in pig-associated strains (Huijsdens *et al.*, 2006), but more studies need to be conducted to confirm this. There are relatively few CA-MRSA strains, with six currently recognized (Eady and Cove, 2003). It has been suggested that these strains are associated with particular niches (Shukla, 2005), concurring with the potential association of the already mentioned sequence types with pigs and pig farmers. The antibiotic resistance patterns of pig MRSA, other than the resistance to methicillin, reflect the antibiotics used in veterinary medicine: tetracycline resistance is very common, whereas ciprofloxacin resistance has been found to be completely absent (de Neeling *et al.*, 2007). This is very different from the situation in nosocomial MRSA strains, in which over 80% of MRSAs may be resistant to ciprofloxacin (Marangon *et al.*, 2004; Raviglione *et al.*, 1990), perhaps suggesting that ST398 is circulating in pigs and only infecting humans occasionally, or that pigs are an emerging reservoir of MRSA.

#### D. Cattle associated MRSA

MRSA is comparatively rare in cows, and those *S. aureus* found to be resistant to methicillin tend to lack SCC*mec*, gaining resistance from the production of  $\beta$ -lactamases (De Oliveira *et al.*, 2000). Penicillin and ampicillin resistance are more widespread (Gentilini *et al.*, 2000; Guler *et al.*, 2005) because of the frequent use of these antibiotics in treating intramammary infections (Pengov and Ceru, 2003). Antibacterial treatment for mastitis has been implicated as a major cause of resistance in bacterial isolates from treated animals, other animals within the herd, and in meat products intended for human consumption (Berghash *et al.*, 1983;

Chen *et al.*, 2004; Griggs *et al.*, 1994; Piddock, 1996). Furthermore, the secretion of antibiotics in milk produced by cows under treatment for mastitis is one of the most common causes of illegal antibacterial residues in milk (Erskine, 1996). Although cattle have a greater freedom of movement than intensively farmed pigs and poultry, effectively reducing the risk of transmission, other risk factors are present. There are extreme inconsistencies in the therapeutic success rates of the standard treatment regimes (Dingwell *et al.*, 2003), with chronic infections being cured through intramammary antibiotic infusions only 35% of the time (Owens *et al.*, 1997). Indiscriminate use of antibiotics, incomplete treatments, and incorrect diagnoses of antibiotic resistant strains do not help the situation (Turutoglu *et al.*, 2006). Although methicillin is not generally used in treatment of cows, MRSA is increasing in its significance as a cause of bovine mastitis (Bernabé *et al.*, 2005; Turutoglu *et al.*, 2006). Acquisition of MRSA may occur through direct contact with humans: one outbreak of mastitis has been attributed to contact with a farm worker (Devriese *et al.*, 1986), and more recent empirical evidence concurs with this (Fox *et al.*, 1991; Roberson *et al.*, 1994). However, this is slightly contentious as evidence to the contrary suggests that transmission between humans and cows is rare (Kapur *et al.*, 1995; Lopes *et al.*, 1990) because of the host specificity of *S. aureus* clones (Smith *et al.*, 2005).

Environmental survival of MRSA may contribute to bovine infections, as shown in the clinical setting as discussed by Hardy *et al.* (2004), where it may survive for several months and act as a source for infecting patients (Neely and Maley, 2000; Sasatsu *et al.*, 1993). Biofilm formation on various materials allows persistence and avoidance of desiccation (Gotz, 2002), so there is a risk of transmission from cow to cow via milking devices and other surfaces. Despite the move towards automated milking practices, reducing contact between humans and untreated milk, the production of aerosols in milking parlors may put workers, as well as other cows, at risk from contamination (Roberson *et al.*, 1994). Similarly, aerial contamination has been noted in pig barns (Gibbs *et al.*, 2004), and in aerosols produced by confined swine-feeding operations (Chapin *et al.*, 2005; Sapkota *et al.*, 2006). Pig feeds have been implicated as potential sources of contamination of tetracycline-resistant *S. aureus*, which may thrive in tetracycline enriched feeds (de Neeling *et al.*, 2007).

## E. Horse associated MRSA

MRSA has also been identified as an emerging disease in horses (Weese *et al.*, 2005a,b, 2006), which might act as a reservoir of rare strains for transmission to humans (Baptiste *et al.*, 2005). The isolation of MRSA from two horses in Canada led to a larger study, which revealed the colonization of 79 hospitalized horses and also of 27 people involved in caring for

these animals, strongly suggestive of frequent transmission between horses and humans (Weese *et al.*, 2005b). All isolates in the study were from nasal swabs and belonged to the Canadian community associated strain C-MRSA5, which is rare in humans. Thus horses may act as a reservoir for this strain, perhaps mirroring the putative association of ST398 with pigs. In Slovenia, MRSA was absent from a total of 300 horses sampled with nasal swabs (Vengust *et al.*, 2006); one explanation for this is that MRSA may also colonize the perineum or the throat, and this has been found to be the sole site of infection in some cases (Coello *et al.*, 1994), so it is possible that subclinical carriers of MRSA were missed. Indeed, a study on humans in nursing homes indicated that up to 14% of cases were missed by screening the nares alone (Lee *et al.*, 1997). MRSA colonization tends to be transient in nature, and so may be missed on screening, whereas it has been shown that infections are likely to persist in hospitalized horses, particularly in those that have undergone invasive procedures (Seguin *et al.*, 1999).

## F. MRSA in companion animals

The situation in companion animals differs from that seen in pigs, as there does not appear to be any association with a particular MRSA clone. The same strains are found in humans and companion animals (van Duijkeren *et al.*, 2003), indicating that pets acquire the infection from humans. Although *S. intermedius* is the predominant staphylococcal species in dogs (Biberstein *et al.*, 1984; Hoekstra and Paulton, 2002), *S. aureus* is also frequently found on the skin or associated with suppurative infections (Kloos, 1990). Dogs may act as a reservoir of MRSA (Baptiste *et al.*, 2005) and have been implicated in causing unexplained relapses in humans previously treated and cleared of MRSA infection (Manian, 2003; van Duijkeren *et al.*, 2003). Strains isolated from humans and dogs have been found to have the same PFGE patterns (Manian, 2003) and SCCmec types (van Duijkeren *et al.*, 2004). Following treatment of dogs, relapses of infection in the owners ceased (Cefai *et al.*, 1994; Manian, 2003; van Duijkeren *et al.*, 2004). Although, like dogs, cats may also carry MRSA, *S. aureus* is not the most common staphylococcal infection; in cats this is fulfilled by a combination of *S. intermedius* and the CoNS *S. felis* (Lilenbaum *et al.*, 1998; Patel *et al.*, 1999). Surgery is a risk factor in cats and dogs just as it is in humans, making up the vast majority of MRSA cases (Boag *et al.*, 2004; Owen *et al.*, 2004; Tomlin *et al.*, 1999; van Duijkeren *et al.*, 2003).

The epidemiology of MRSA appears to be changing, with the possibility of animals acting as reservoirs of infection for people without any of the usual risk factors. In countries that currently have a low incidence of MRSA, this may be an important source of an otherwise rare infection.

## VI. CONCLUSIONS

The evolution of antibiotic resistant bacteria is one of the most significant problems in modern medicine and poses a serious threat to human health. Increasingly the huge diversity of resistance genes that already exist in the environment is beginning to be appreciated. Understanding the selective pressures and mechanisms of gene transfer that drive dissemination of resistance genes not only in the clinic, but also in the wider environment is crucial for long-term strategies in the treatment of microbial disease.

Modern farming practice is attempting to reduce dependency on antibiotics but this in itself may not reduce particular mechanisms of resistance such as genes carried on class 1 integrons, which we have shown can be selected for by biocides in the environment. Understanding the ecology of resistance genes is extremely difficult as genes may be carried by unculturable bacteria (99.0–99.9% of bacteria). Movement of genes between environmental bacteria and the clinic has therefore been difficult to investigate in the past. However, modern molecular approaches such as epidemiological studies of key resistance determinants in total community DNA using quantitative real-time PCR allows detailed analyses and comparison of gene prevalence in the environment and human gut. Functional metagenomics and integron clone library construction allows the entire resistance gene pool or metagenome to be analyzed. Antibiotic resistant bacteria in the environment may be transferred to the human population via ingestion of contaminated food and water, via direct contact with animals, swimming in lakes, rivers, and the sea, and by airborne bacteria. In reality, there is no distinction between clinical and nonclinical environments; both must be considered to fully understand the underlying causes of antibiotic resistance in clinically important bacteria.

## REFERENCES

- Aarestrup, F. M. (2000). Characterization of glycopeptide-resistant enterococcus faecium (GRE) from broilers and pigs in Denmark: Genetic evidence that persistence of GRE in pig herds is associated with coselection by resistance to macrolides. *J. Clin. Microbiol.* **38**, 2774–2777.
- Abraham, E. P., and Chain, E. (1940). An enzyme from bacteria able to destroy penicillin. *Nature* **146**, 837.
- Agerso, Y., Sengelov, G., and Jensen, L. B. (2004). Development of a rapid method for direct detection of *tet(M)* genes in soil from Danish farmland. *Environ. Int.* **30**, 117–122.
- Agustía, C. M., Pujola, M., Argerichb, M. J., Ayatsc, J., Badiad, M., Domínguezc, M. A., Corbellaa, X., and Arizaa, J. (2002). Short-term effect of the application of selective decontamination of the digestive tract on different body site reservoir ICU

- patients colonized by multi-resistant *Acinetobacter baumannii*. *J. Antimicrob. Chemother.* **49**, 205–208.
- Anthonisen, I. L., Sunde, M., Steinum, T. M., Sidhu, M. S., and Sorum, H. (2002). Organization of the antiseptic resistance Gene *qacA* and Tn552-related  $\beta$ -lactamase genes in multidrug-resistant *Staphylococcus haemolyticus* strains of animal and human origins. *Antimicrob. Agents Chemother.* **46**, 3606–3612.
- Archer, G. L., and Niemerger, D. M. (1994). Origin and evolution of DNA associated with resistance to methicillin in staphylococci. *Trends Microbiol.* **2**, 343–347.
- Armand-Lefevre, L., Ruimy, R., and Andremont, A. (2005). Clonal comparison of *Staphylococcus aureus* isolates from healthy pig farmers, human controls, and pigs. *Emerg. Infect. Dis.* **11**, 711–714.
- Asensio, A., Guerrero, A., Quereda, C., Lizan, M., and Martinez-Ferrer, M. (1996). Colonization and infection with methicillin-resistant *Staphylococcus aureus*: Associated factors and eradication. *Infect. Control Hosp. Epidemiol.* **17**, 20–28.
- Aubry-Damon, H., Grenet, K., Sall-Ndiaye, P., Che, D., Cordeiro, E., Bougnoux, M. E., Rigaud, E., Le Strat, Y., Lemanissier, V., Armand-Lefevre, L., Delzescaux, D., Desenclos, J. C., et al. (2004). Antimicrobial resistance in commensal flora of pig farmers. *Emerg. Infect. Dis.* **10**, 873–879.
- Baptiste, K. E., Williams, K., Willams, N. J., Wattret, A., Clegg, P. D., Dawson, S., Corkill, J. E., O'Neill, T., and Hart, C. A. (2005). Methicillin-resistant staphylococci in companion animals. *Emerg. Infect. Dis.* **11**, 1942–1944.
- Bass, L., Liebert, C. A., Lee, M. D., Summers, A. O., White, D. G., Thayer, S. G., and Maurer, J. J. (1999). Incidence and characterization of integrons, genetic elements mediating multiple-drug resistance, in avian *Escherichia coli*. *Antimicrob. Agents Chemother.* **43**, 2925–2929.
- Berg, G., Eberl, L., and Hartmann, A. (2005). The rhizosphere as a reservoir for opportunistic human pathogenic bacteria. *Environ. Microbiol.* **7**, 1673–1685.
- Berghash, S. R., Davidson, J. N., Armstrong, J. C., and Dunny, G. M. (1983). Effects of antibiotic treatment of nonlactating dairy cows on antibiotic resistance patterns of bovine mastitis pathogens. *Antimicrob. Agents Chemother.* **24**, 771–776.
- Bernabé, L., Ordóñez, V., and Vázquez, J. C. (2005). Identification in cows presenting subclinical mastitis. *Animals and environment, Vol. 1: Proceedings of the XIIth ISAH Congress on Animal Hygiene*, pp. 308–310. Warsaw, Poland, September 2005, 4–8.
- Biberstein, E. L., Jang, S. S., and Hirsh, D. C. (1984). Species distribution of coagulase-positive staphylococci in animals. *J. Clin. Microbiol.* **19**, 610–615.
- Boag, A., Loeffler, A., and Lloyd, D. H. (2004). Methicillin-resistant *Staphylococcus aureus* isolates from companion animals. *Vet. Rec.* **154**, 411.
- Bonnet, R. (2004). Growing group of extended-spectrum beta-lactamases: The CTX-M enzymes. *Antimicrob. Agents Chemother.* **48**, 1–14.
- Boxall, A. B., Blackwell, P., Cavallo, R., Kay, P., and Tolls, J. (2002). The sorption and transport of a sulphonamide antibiotic in soil systems. *Toxicol. Lett.* **131**, 19–28.
- Boyle-Vavra, S., and Daum, R. S. (2007). Community-acquired methicillin-resistant *Staphylococcus aureus*: The role of Panton–Valentine leukocidin. *Lab. Invest.* **87**, 3–9.
- Briggs, C. E., and Fratamico, P. M. (1999). Molecular characterization of an antibiotic resistance gene cluster of *Salmonella typhimurium* DT104. *Antimicrob. Agents Chemother.* **43**, 846–849.
- Brinas, L., Moreno, M. A., Teshager, T., Zarazaga, M., Saenz, Y., Porrero, C., Dominguez, L., and Torres, C. (2003a). Beta-lactamase characterization in *Escherichia coli* isolates with diminished susceptibility or resistance to extended-spectrum cephalosporins recovered from sick animals in Spain. *Microb. Drug Resist.* **9**, 201–209.
- Brinas, L., Moreno, M. A., Zarazaga, M., Porrero, C., Saenz, Y., Garcia, M., Dominguez, L., and Torres, C. (2003b). Detection of CMY-2, CTX-M-14, and SHV-12 beta-lactamases in

- Escherichia coli* fecal-sample isolates from healthy chickens. *Antimicrob. Agents Chemother.* **47**, 2056–2058.
- Brinas, L., Zarazaga, M., Saenz, Y., Ruiz-Larrea, F., and Torres, C. (2002). Beta-lactamases in ampicillin-resistant *Escherichia coli* isolates from foods, humans, and healthy animals. *Antimicrob. Agents Chemother.* **46**, 3156–3163.
- Burd, G. I., Dixon, D. G., and Glick, B. R. (1998). A plant growth-promoting bacterium that decreases nickel toxicity in seedlings. *Appl. Environ. Microbiol.* **64**, 3663–3668.
- Burd, G. I., Dixon, D. G., and Glick, B. R. (2000). Plant growth-promoting bacteria that decrease heavy metal toxicity in plants. *Can. J. Microbiol.* **46**, 237–245.
- Burrus, V., and Waldor, M. K. (2004). Shaping bacterial genomes with integrative and conjugative elements. *Res. Microbiol.* **155**, 376–386.
- Busato, A., Trachsel, P., Schallibaum, M., and Blum, J. W. (2000). Udder health and risk factors for subclinical mastitis in organic dairy farms in Switzerland. *Prev. Vet. Med.* **44**, 205–220.
- Cefai, C., Ashurst, S., and Owens, C. (1994). Human carriage of methicillin-resistant *Staphylococcus aureus* linked with pet dog. *Lancet* **344**, 539–540.
- Chapin, A., Rule, A., Gibson, K., Buckley, T., and Schwab, K. (2005). Airborne multidrug-resistant bacteria isolated from a concentrated swine feeding operation. *Environ. Health Perspect.* **113**, 137–142.
- Chee-Sanford, J. C., Aminov, R. I., Krapac, I. J., Garrigues-Jeanjean, N., and Mackie, R. I. (2001). Occurrence and diversity of tetracycline resistance genes in lagoons and ground-water underlying two swine production facilities. *Appl. Environ. Microbiol.* **67**, 1494–1502.
- Chen, S., Zhao, S., White, D. G., Schroeder, C. M., Lu, R., Yang, H., McDermott, P. F., Ayers, S., and Meng, J. (2004). Characterization of multiple-antimicrobial-resistant *Salmonella* Serovars Isolated from Retail Meats. *Appl. Environ. Microbiol.* **70**, 1–7.
- Coello, R., Jimenez, J., Garcia, M., Arroyo, P., Minguez, D., Fernandez, C., Cruzet, F., and Gaspar, C. (1994). Prospective study of infection, colonization and carriage of methicillin-resistant *Staphylococcus aureus* in an outbreak affecting 990 patients. *Eur. J. Clin. Microbiol. Infect. Dis.* **13**, 74–81.
- Coffey, T. J., Dowson, C. G., Daniels, M., and Spratt, B. G. (1995). Genetics and molecular biology of beta-lactam-resistant pneumococci. *Microb. Drug Resist.* **1**, 29–34.
- Couto, I., de Lencastre, H., Severina, E., Kloos, W., Webster, J. A., Hubner, R. J., Sanches, I. S., and Tomasz, A. (1996). Ubiquitous presence of a meca homologue in natural isolates of *Staphylococcus sciuri*. *Microb. Drug Resist. (Larchmont, N.Y.)* **2**, 377–391.
- Couto, I., Wu, S. W., Tomasz, A., and de Lencastre, H. (2003). Development of methicillin resistance in clinical isolates of *Staphylococcus sciuri* by transcriptional activation of the meca homologue native to the species. *J. Bacteriol.* **185**, 645–653.
- Cove, J. H., Eady, E. A., and Cunliffe, W. J. (1990). Skin carriage of antibiotic-resistant coagulase-negative staphylococci in untreated subjects. *J. Antimicrob. Chemother.* **25**, 459–469.
- Dar, J. A., Thoker, M. A., Khan, J. A., Ali, A., Khan, M. A., Rizwan, M., Bhat, K. H., Dar, M. J., Ahmed, N., and Ahmad, S. (2006). Molecular epidemiology of clinical and carrier strains of methicillin resistant *Staphylococcus aureus* (MRSA) in the hospital settings of north India. *Ann. Clin. Microbiol. Antimicrob.* **5**, 22.
- Daum, R. S., Ito, T., Hiramatsu, K., Hussain, F., Mongkolrattanothai, K., Jamklang, M., and Boyle-Vavra, S. (2002). A novel methicillin-resistance cassette in community-acquired methicillin-resistant *Staphylococcus aureus* isolates of diverse genetic backgrounds. *J. Infect. Dis.* **186**, 1344–1347.
- Davidson, J. (1999). Genetic exchange between bacteria in the environment. *Plasmid* **42**, 73–91.
- De Liguoro, M., Cibir, V., Capolongo, F., Halling-Sorensen, B., and Montesissa, C. (2003). Use of oxytetracycline and tylosin in intensive calf farming: Evaluation of transfer to manure and soil. *Chemosphere* **52**, 203–212.
- de Neeling, A. J., van den Broek, M. J., Spalburg, E. C., van Santen-Verheuevel, M. G., Dam-Deisz, W. D., Boshuizen, H. C., van de Giessen, A. W., van Duijkeren, E., and



- Huijsdens, X. W. (2007). High prevalence of methicillin resistant *Staphylococcus aureus* in pigs. *Vet. Microbiol.* **122**, 366–372.
- De Oliveira, A. P., Watts, J. L., Salmon, S. A., and Aarestrup, F. M. (2000). Antimicrobial susceptibility of *Staphylococcus aureus* isolated from bovine mastitis in Europe and the United States. *J. Dairy Sci.* **83**, 855–862.
- Decousser, J. W., Poirel, L., and Nordmann, P. (2001). Characterization of a chromosomally encoded extended-spectrum class A beta-lactamase from *Kluyvera cryocrescens*. *Antimicrob. Agents Chemother.* **45**, 3595–3598.
- Devriese, L. A., Hommez, J., Kilpper-Baelz, R., and Schleifer, K. H. (1986). *Streptococcus canis* sp. Nov.: A species of group G streptococci from animals. *Int. J. Syst. Bacteriol.* **36**, 422–425.
- Dingwell, R. T., Leslie, K. E., Duffield, T. F., Schukken, Y. H., DesCoteaux, L., Keefe, G. P., Kelton, D. F., Lissemore, K. D., Shewfelt, W., Dick, P., and Bagg, R. (2003). Efficacy of intramammary tilmicosin and risk factors for cure of *Staphylococcus aureus* infection in the dry period. *J. Dairy Sci.* **86**, 159–168.
- Dunny, G. M., Antiporta, M. H., and Hirt, H. (2001). Peptide pheromone-induced transfer of plasmid pcf10 in *Enterococcus faecalis*: Probing the genetic and molecular basis for specificity of the pheromone response. *Peptides* **22**, 1529–1539.
- Eady, E. A., and Cove, J. H. (2003). Staphylococcal resistance revisited: Community-acquired methicillin resistant *Staphylococcus aureus*—An emerging problem for the management of skin and soft tissue infections. *Curr. Opin. Infect. Dis.* **16**, 103–124.
- Ensor, V., Warren, R., Neill, P. O., Butler, V., Taylor, J., Nye, K., Harvey, M., Livermore, D., Woodford, N., and Hawkey, P. (2007). Isolation of quinolone-resistance CTX-M-producing *Escherichia coli* from raw chicken meat sold in retail outlets in the West Midlands, U.K. In “17th European Congress of Clinical Microbiology and Infectious Diseases,” pp. poster number 1025, Munich.
- Erskine, R. J. (1996). Why do antibiotic residues in milk happen. *Michigan Dairy Rev.* **1**, 16.
- Fanaro, S., Chierici, R., Guerrini, P., and Vigi, V. (2003). Intestinal microflora in early infancy: Composition and development. *Acta Paediatr. Suppl.* **91**, 48–55.
- Fluit, A. C., and Schmitz, F. J. (2004). Resistance integrons and super-integrons. *Clin. Microbiol. Infect.* **10**, 272–288.
- Fox, L. K., Gershman, M., Hancock, D. D., and Hutton, C. T. (1991). Fomites and reservoirs of *Staphylococcus aureus* causing intramammary infections as determined by phage typing: The effect of milking time hygiene practices. *Cornell Vet.* **81**, 183–193.
- Freter, R., Freter, R. R., and Brickner, H. (1983). Experimental and mathematical models of *Escherichia coli* plasmid transfer *in vitro* and *in vivo*. *Infect. Immun.* **39**, 60–84.
- Gaze, W. H., Abdoulsam, N., Hawkey, P. M., and Wellington, E. M. (2005). Incidence of class 1 integrons in a quaternary ammonium compound-polluted environment. *Antimicrob. Agents Chemother.* **49**, 1802–1807.
- Gentilini, E., Denamiel, G., Llorente, P., Godaly, S., Rebuelto, M., and DeGregorio, O. (2000). Antimicrobial susceptibility of *Staphylococcus aureus* isolated from bovine mastitis in Argentina. *J. Dairy Sci.* **83**, 1224–1227.
- Germida, J., and Siciliano, S. (2001). Taxonomic diversity of bacteria associated with the roots of modern, recent and ancient wheat cultivars. *Biol. Fertil. Soils* **33**, 410–415.
- Gibbs, S. G., Green, C. F., Tarwater, P. M., and Scarpino, P. V. (2004). Airborne antibiotic resistant and nonresistant bacteria and fungi recovered from two swine herd confined animal feeding operations. *J. Occup. Environ. Hyg.* **1**, 699–706.
- Gill, S. R., Fouts, D. E., Archer, G. L., Mongodin, E. F., Deboy, R. T., Ravel, J., Paulsen, I. T., Kolonay, J. F., Brinkac, L., Beanan, M., Dodson, R. J., and Daugherty, S. C. (2005). Insights on evolution of virulence and resistance from the complete genome analysis of an early methicillin-resistant *Staphylococcus aureus* strain and a biofilm-producing methicillin-resistant *Staphylococcus epidermidis* strain. *J. Bacteriol.* **187**, 2426–2438.

- Golet, E. M., Xifra, I., Siegrist, H., Alder, A. C., and Giger, W. (2003). Environmental exposure assessment of fluoroquinolone antibacterial agents from sewage to soil. *Environ. Sci. Technol.* **37**, 3243–3249.
- Gotz, F. (2002). *Staphylococcus* and biofilms. *Mol. Microbiol.* **43**, 1367–1378.
- Griggs, D. J., Hall, M. C., Jin, Y. F., and Piddock, L. J. (1994). Quinolone resistance in veterinary isolates of *Salmonella*. *J. Antimicrob. Chemother.* **33**, 1173–1189.
- Guler, L., Ok, U., Gunduz, K., Gulcu, Y., and Hadimli, H. H. (2005). Antimicrobial susceptibility and coagulase gene typing of *Staphylococcus aureus* isolated from bovine clinical mastitis cases in Turkey. *J. Dairy Sci.* **88**, 3149–3154.
- Halbert, L. W., Kaneene, J. B., Ruegg, P. L., Warnick, L. D., Wells, S. J., Mansfield, L. S., Fossler, C. P., Campbell, A. M., and Geiger-Zwald, A. M. (2006). Evaluation of antimicrobial susceptibility patterns in *Campylobacter* spp. isolated from dairy cattle and farms managed organically and conventionally in the midwestern and northeastern United States. *J. Am. Vet. Med. Assoc.* **228**, 1074–1081.
- Hall, R. M., Brookes, D. E., and Stokes, H. W. (1991). Site-specific insertion of genes into integrons: Role of the 59-base element and determination of the recombination cross-over point. *Mol. Microbiol.* **5**, 1941–1959.
- Hamscher, G., Sczesny, S., Hoper, H., and Nau, H. (2002). Determination of persistent tetracycline residues in soil fertilized with liquid manure by high-performance liquid chromatography with electrospray ionization tandem mass spectrometry. *Anal. Chem.* **74**, 1509–1518.
- Hansen-Nord, M., Gahrn-Hansen, B., and Siboni, K. (1988). Studies on clinical isolates of coagulase-negative staphylococci resistant to methicillin. Evidence of cross-resistance between methicillin and cephalothin. *APMIS* **96**, 133–140.
- Hanssen, A. M., and Ericson Sollid, J. U. (2006). SCCmec in staphylococci: Genes on the move. *FEMS Immunol. Med. Microbiol.* **46**, 8–20.
- Hanssen, A. M., Kjeldsen, G., and Sollid, J. U. E. (2004). Local variants of Staphylococcal cassette chromosome *mec* in sporadic methicillin-resistant *Staphylococcus aureus* and methicillin-resistant coagulase-negative Staphylococci: Evidence of horizontal gene transfer? *Antimicrob. Chemother.* **48**, 285–296.
- Hardy, K. J., Hawkey, P. M., Gao, F., and Oppenheim, B. A. (2004). Methicillin resistant *Staphylococcus aureus* in the critically ill. *Br. J. Anaesth.* **92**, 121–130.
- Hartman, B., and Tomasz, A. (1981). Altered penicillin-binding proteins in methicillin-resistant strains of *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* **19**, 726–735.
- Hasman, H., Kempf, I., Chidaine, B., Cariolet, R., Ersboll, A. K., Houe, H., Bruun Hansen, H. C., and Aarestrup, F. M. (2006). Copper resistance in *Enterococcus faecium*, mediated by the *trb* gene, is selected by supplementation of pig feed with copper sulfate. *Appl. Environ. Microbiol.* **72**, 5784–5789.
- Hawes, J. (2004). The impact of and resistance to fluoroquinolones in the environment. In "Biology," p. 252. University of Warwick, England.
- Hawkey, P. M. (1998). The origins and molecular basis of antibiotic resistance. *BMJ* **317**, 657–660.
- Henriques, I., Moura, A., Alves, A., Saavedra, M. J., and Correia, A. (2006). Analysing diversity among beta-lactamase encoding genes in aquatic environments. *FEMS Microbiol. Ecol.* **56**, 418–429.
- Herold, B. C., Immergluck, L. C., Maranan, M. C., Lauderdale, D. S., Gaskin, R. E., Boyle-Vavra, S., Leitch, C. D., and Daum, R. S. (1998). Community-acquired methicillin-resistant *Staphylococcus aureus* in children with no identified predisposing risk. *JAMA* **279**, 593–598.
- Heuer, O. E., Pedersen, K., Andersen, J. S., and Madsen, M. (2002a). Vancomycin-resistant enterococci (VRE) in broiler flocks 5 years after the avoparcin ban. *Microb. Drug Resist.* **8**, 133–138.

- Heuer, O. E., Pedersen, K., Jensen, L. B., Madsen, M., and Olsen, J. E. (2002b). Persistence of vancomycin-resistant enterococci (VRE) in broiler houses after the avoparcin ban. *Microb. Drug Resist.* **8**, 355–361.
- Hiramatsu, K., Cui, L., Kuroda, M., and Ito, T. (2001). The emergence and evolution of methicillin-resistant *Staphylococcus aureus*. *Trends Microbiol.* **9**, 486–493.
- Hoekstra, K. A., and Paulton, R. J. (2002). Clinical prevalence and antimicrobial susceptibility of *Staphylococcus aureus* and *Staph. intermedius* in dogs. *J. Appl. Microbiol.* **93**, 406–413.
- Hollander, R., Ebke, M., Barck, H., and von Pritzbuere, E. (2001). Asymptomatic carriage of *Klebsiella pneumoniae* producing extended-spectrum beta-lactamase by patients in a neurological early rehabilitation unit: Management of an outbreak. *J. Hosp. Infect.* **48**, 207–213.
- Holmes, A. J., Gillings, M. R., Nield, B. S., Mabbutt, B. C., Nevalainen, K. M., and Stokes, H. W. (2003). The gene cassette metagenome is a basic resource for bacterial genome evolution. *Environ. Microbiol.* **5**, 383–394.
- Huijsdens, X. W., van Dijke, B. J., Spalburg, E., van Santen-Verheuevel, M. G., Heck, M. E. O. C., Pluister, G. N., Voss, A., Wannet, W. J. B., and de Neeling, A. J. (2006). Community-acquired MRSA and pig-farming. *Ann. Clin. Microbiol. Antimicrob.* **5**, 26.
- Hutchison, M. L., Walters, L. D., Moore, A., Crookes, K. M., and Avery, S. M. (2004). Effect of length of time before incorporation on survival of pathogenic bacteria present in livestock wastes applied to agricultural soil. *Appl. Environ. Microbiol.* **70**, 5111–5118.
- Ito, T., Katayama, Y., and Hiramatsu, K. (1999). Cloning and nucleotide sequence determination of the entire *mec* DNA of pre-methicillin-resistant *Staphylococcus aureus* N315. *Antimicrob. Agents Chemother.* **43**, 1449–1458.
- Ito, T., Okuma, K., Ma, X. X., Yuzawa, H., and Hiramatsu, K. (2003). Insights on antibiotic resistance of *Staphylococcus aureus* from its whole genome: Genomic island SCC. *Drug Resist. Updat.* **6**, 41–52.
- Jensen, S. E., and Demain, A. L. (1995). Beta-lactams. *Biotechnology* **28**, 239–268.
- Kapur, V., Sischo, W. M., Greer, R. S., Whittam, T. S., and Musser, J. M. (1995). Molecular population genetic analysis of *Staphylococcus aureus* recovered from cows. *J. Clin. Microbiol.* **33**, 376–380.
- Katayama, Y., Zhang, H. Z., Hong, D., and Chambers, H. F. (2003). Jumping the Barrier to  $\beta$ -lactam Resistance in *Staphylococcus aureus*. *J. Bacteriol.* **185**, 5465–5472.
- Kirby, W. M. M. (1944). Extraction of a highly potent penicillin inactivator from penicillin resistant staphylococci. *Science* 452–453.
- Kloos, W. E. (1990). Systematics and the natural history of staphylococci 1. *Soc. Appl. Bacteriol. Symp. Ser.* **19**, 25–37.
- Kummerer, K. (2004). Resistance in the environment. *J. Antimicrob. Chemother.* **54**, 311–320.
- Kuroda, M., Ohta, T., Uchiyama, I., Baba, T., Yuzawa, H., Kobayashi, I., Cui, L., Oguchi, A., Aoki, K., and Nagai, Y. (2001). Whole genome sequencing of methicillin-resistant *Staphylococcus aureus*. *Lancet* **357**, 1225–1240.
- Lee, Y. J., Cho, J. K., Kim, K. S., Tak, R. B., Kim, A. R., Kim, J. W., Im, S. K., and Kim, B. H. (2005). Fluoroquinolone resistance and *gyrA* and *parC* mutations of *Escherichia coli* isolated from chicken. *J. Microbiol.* **43**, 391–397.
- Lee, Y. L., Cesario, T., Gupta, G., Flionis, L., Tran, C., Decker, M., and Thrupp, L. (1997). Surveillance of colonization and infection with *Staphylococcus aureus* susceptible or resistant to methicillin in a community skilled-nursing facility. *Am. J. Infect. Control* **25**, 312–321.
- Leverstein-van Hall, M. A., He, M. B., Ar, T. D., Paaeuw, A., Fluit, A. C., and Verhoef, J. (2003). Multidrug resistance among Enterobacteriaceae is strongly associated with the presence of integrons and is independent of species or isolate origin. *J. Infect. Dis.* **187**, 251–259.
- Levy, S. B., FitzGerald, G. B., and Macone, A. B. (1976). Changes in intestinal flora of farm personnel after introduction of a tetracycline-supplemented feed on a farm. *N. Engl. J. Med.* **295**, 583–588.

- Levy, S. B., Marshall, B., Schluenderberg, S., Rowse, D., and Davis, J. (1988). High frequency of antimicrobial resistance in human fecal flora. *Antimicrob. Agents Chemother.* **32**, 1801–1806.
- Lilenbaum, W., Nunes, E. L., and Azeredo, M. A. (1998). Prevalence and antimicrobial susceptibility of staphylococci isolated from the skin surface of clinically normal cats. *Lett. Appl. Microbiol.* **27**, 224–228.
- Lindsay, J. A., and Holden, M. T. (2004). *Staphylococcus aureus*: Superbug, super genome? *Trends Microbiol.* **12**, 378–385.
- Livermore, D. M., and Hawkey, P. M. (2005). CTX-M: changing the face of esbls in the UK. *J. Antimicrob. Chemother.* **56**, 451–454.
- Lopes, C. A., Moreno, G., Curi, P. R., Gottschalk, A. F., Modolo, J. R., Horacio, A., Correa, A., and Pavan, C. (1990). Characteristics of *Staphylococcus aureus* from subclinical bovine mastitis in Brazil. *Br. Vet. J.* **146**, 443–448.
- Machado, E., Coque, T. M., Canton, R., Baquero, F., Sousa, J. C., and Peixe, L. (2006). Dissemination in Portugal of CTX-M-15-, OXA-1-, and TEM-1-producing *Enterobacteriaceae* strains containing the *aac(6′)-Ib-cr* gene, which encodes an aminoglycoside- and fluoroquinolone-modifying enzyme. *Antimicrob. Agents Chemother.* **50**, 3220–3221.
- Mahamoud, A., Chevalier, J., Alibert-Franco, S., Kern, W. V., and Pages, J. M. (2007). Antibiotic efflux pumps in Gram-negative bacteria: The inhibitor response strategy. *J. Antimicrob. Chemother.* **59**, 1223–1229.
- Manian, F. A. (2003). Asymptomatic nasal carriage of mupirocin-resistant, methicillin-resistant *Staphylococcus aureus* (MRSA) in a pet dog associated with MRSA infection in household contacts. *Clin. Infect. Dis.* **36**, 26–28.
- Marangon, F. B., Miller, D., Muallem, M. S., Romano, A. C., and Alfonso, E. C. (2004). Ciprofloxacin and levofloxacin resistance among methicillin-sensitive *Staphylococcus aureus* isolates from keratitis and conjunctivitis. *Am. J. Ophthalmol.* **137**, 453–458.
- McDonnell, R. W., Sweeney, H. M., and Cohen, S. (1983). Conjugational transfer of gentamicin resistance plasmids intra- and interspecifically in *Staphylococcus aureus* and *Staphylococcus epidermidis*. *Antimicrob. Agents Chemother.* **23**, 151–160.
- Menzies, R. E., Cornere, B. M., and MacCulloch, D. (1987). Cephalosporin susceptibility of methicillin-resistant, coagulase-negative staphylococci. *Antimicrob. Agents Chemother.* **31**, 42–45.
- Mesa, R. J., Blanc, V., Blanch, A. R., Cortes, P., Gonzalez, J. J., Lavilla, S., Miro, E., Muniesa, M., Saco, M., Tortola, M. T., and Mirelis, B. (2006). Extended-spectrum beta-lactamase-producing *Enterobacteriaceae* in different environments (humans, food, animal farms and sewage). *J. Antimicrob. Chemother.* **58**, 211–215.
- Morales, A., Garland, J. L., and Lim, D. V. (1996). Survival of potentially pathogenic human-associated bacteria in the rhizosphere of hydroponically grown wheat. *FEMS Microbiol. Ecol.* **20**, 155–162.
- Nandi, S., Maurer, J. J., Hofacre, C., and Summers, A. O. (2004). Gram-positive bacteria are a major reservoir of Class 1 antibiotic resistance integrons in poultry litter. *Proc. Natl. Acad. Sci. USA.* **101**, 7118–7122.
- Neely, A. N., and Maley, M. P. (2000). Survival of enterococci and staphylococci on hospital fabrics and plastic. *J. Clin. Microbiol.* **38**, 724–726.
- Nemergut, D. R., Martin, A. P., and Schmidt, S. K. (2004). Integron diversity in heavy-metal-contaminated mine tailings and inferences about integron evolution. *Appl. Environ. Microbiol.* **70**, 1160–1168.
- Nield, B. S., Holmes, A. J., Gillings, M. R., Recchia, G. D., Mabbutt, B. C., Nevalainen, K. M., and Stokes, H. W. (2001). Recovery of new integron classes from environmental DNA. *FEMS Microbiol. Lett.* **195**, 59–65.
- Nordmann, P., and Poirel, L. (2005). Emergence of plasmid-mediated resistance to quinolones in *Enterobacteriaceae*. *J. Antimicrob. Chemother.* **56**, 463–469.

- Owen, M. R., Moores, A. P., and Coe, R. J. (2004). Management of MRSA septic arthritis in a dog using a gentamicin-impregnated collagen sponge. *J. Small Anim. Pract.* **45**, 609–612.
- Owens, W. E., Ray, C. H., Watts, J. L., and Yancey, R. J. (1997). Comparison of success of antibiotic therapy during lactation and results of antimicrobial susceptibility tests for bovine mastitis. *J. Dairy Sci.* **80**, 313–317.
- Partridge, S. R., Brown, H. J., and Hall, R. M. (2002). Characterization and movement of the class 1 integron known as Tn2521 and Tn1405. *Antimicrob. Agents Chemother.* **46**, 1288–1294.
- Patel, A., Lloyd, D. H., and Lampion, D. S. (1999). Short communication. Antimicrobial resistance of feline staphylococci in south-eastern England. *Vet. Dermatol.* **10**, 257–261.
- Paulsen, I. T., Brown, M. H., and Skurray, R. A. (1996). Proton-dependent multidrug efflux systems. *Microbiol. Rev.* **60**, 575–608.
- Peacock, S. J., de Silva, I., and Lowy, F. D. (2001). What determines nasal carriage of *Staphylococcus aureus*? *Trends Microbiol.* **9**, 605–610.
- Pengov, A., and Ceru, S. (2003). Antimicrobial drug susceptibility of *Staphylococcus aureus* strains isolated from bovine and ovine mammary glands. *J. Dairy Sci.* **86**, 3157–3163.
- Piddock, L. J. (1996). Does the use of antimicrobial agents in veterinary medicine and animal husbandry select antibiotic-resistant bacteria that infect man and compromise antimicrobial chemotherapy?. *J. Antimicrob. Chemother.* **38**, 1–3.
- Piddock, L. J. (1998). Fluoroquinolone resistance: Overuse of fluoroquinolones in human and veterinary medicine can breed resistance. *BMJ* **317**, 1029–1030.
- Pirnay, J. P., De Vos, D., Mossialos, D., Vanderkelen, A., Cornelis, P., and Zizi, M. (2002). Analysis of the *Pseudomonas aeruginosa oprd* gene from clinical and environmental isolates. *Environ. Microbiol.* **4**, 872–882.
- Poirel, L., Liard, A., Rodriguez-Martinez, J. M., and Nordmann, P. (2005a). Vibrionaceae as a possible source of Qnr-like quinolone resistance determinants. *J. Antimicrob. Chemother.* **56**, 1118–1121.
- Poirel, L., Naas, T., Le Thomas, I., Karim, A., Bingen, E., and Nordmann, P. (2001). CTX-M-type extended-spectrum beta-lactamase that hydrolyzes ceftazidime through a single amino acid substitution in the omega loop. *Antimicrob. Agents Chemother.* **45**, 3355–3361.
- Poirel, L., Rodriguez-Martinez, J. M., Mammeri, H., Liard, A., and Nordmann, P. (2005b). Origin of plasmid-mediated quinolone resistance determinant qnrA. *Antimicrob. Agents Chemother.* **49**, 3523–3525.
- Poirel, L., Van De Loo, M., Mammeri, H., and Nordmann, P. (2005c). Association of plasmid-mediated quinolone resistance with extended-spectrum beta-lactamase VEB-1. *Antimicrob. Agents Chemother.* **49**, 3091–3094.
- Rahme, L. G., Stevens, E. J., Wolfort, S. F., Shao, J., Tompkins, R. G., and Ausubel, F. M. (1995). Common virulence factors for bacterial pathogenicity in plants and animals. *Science* **268**, 1899–1902.
- Raviglione, M. C., Boyle, J. F., Mariuz, P., Pablos-Mendez, A., Cortes, H., and Merlo, A. (1990). Ciprofloxacin-resistant methicillin-resistant *Staphylococcus aureus* in an acute-care hospital. *Antimicrob. Agents Chemother.* **34**, 2050–2054.
- Recchia, G. D., and Hall, R. M. (1995). Gene cassettes: A new class of mobile element. *Microbiology* **141**(Pt 12), 3015–3027.
- Roberson, J. R., Fox, L. K., Hancock, D. D., Gay, J. M., and Besser, T. E. (1994). Ecology of *Staphylococcus aureus* isolated from various sites on dairy farms. *J. Dairy Sci.* **77**, 3354–3364.
- Robicsek, A., Jacoby, G. A., and Hooper, D. C. (2006). The worldwide emergence of plasmid-mediated quinolone resistance. *Lancet Infect. Dis.* **6**, 629–640.
- Rodriguez, M. M., Power, P., Radice, M., Vay, C., Famiglietti, A., Galleni, M., Ayala, J. A., and Gutkind, G. (2004). Chromosome-encoded CTX-M-3 from *Kluyvera ascorbata*: A possible origin of plasmid-borne CTX-M-1-derived cefotaximases. *Antimicrob. Agents Chemother.* **48**, 4895–4897.

- Sapkota, A. R., Ojo, K. K., Roberts, M. C., and Schwab, K. J. (2006). Antibiotic resistance genes in multidrug-resistant *Enterococcus* spp. and *Streptococcus* spp. recovered from the indoor air of a large-scale swine-feeding operation. *Lett. Appl. Microbiol.* **43**, 534–540.
- Sasatsu, M., Shibata, Y., Noguchi, N., Nagata, M., Murata, M., Ogata, Y., and Kono, M. (1993). Survival ability of clinical isolates of methicillin-resistant *Staphylococcus aureus*. *Microbios* **75**, 17–21.
- Sato, K., Bartlett, P. C., Kaneene, J. B., and Downes, F. P. (2004). Comparison of prevalence and antimicrobial susceptibilities of *Campylobacter* spp. isolates from organic and conventional dairy herds in Wisconsin. *Appl. Environ. Microbiol.* **70**, 1442.
- Sato, K., Bartlett, P. C., and Saeed, M. A. (2005). Antimicrobial susceptibility of *Escherichia coli* isolates from dairy farms using organic versus conventional production methods. *J. Am. Vet. Med. Assoc.* **226**, 589–594.
- Segal, H., Thomas, R., and Gay Elisha, B. (2003). Characterization of class 1 integron resistance gene cassettes and the identification of a novel IS-like element in *Acinetobacter baumannii*. *Plasmid* **49**, 169–178.
- Seguin, J. C., Walker, R. D., Caron, J. P., Kloos, W. E., George, C. G., Hollis, R. J., Jones, R. N., and Pfaller, M. A. (1999). Methicillin-resistant *Staphylococcus aureus* outbreak in a veterinary teaching hospital: Potential human-to-animal transmission. *J. Clin. Microbiol.* **37**, 1459–1463.
- Shea, K. M. (2004). Nontherapeutic use of antimicrobial agents in animal agriculture: implications for pediatrics. *Pediatrics* **114**, 862–868.
- Shiraki, Y., Shibata, N., Doi, Y., and Arakawa, Y. (2004). *Escherichia coli* producing CTX-M-2 beta-lactamase in cattle, Japan. *Emerg. Infect. Dis.* **10**, 69–75.
- Shukla, S. K. (2005). Community-associated methicillin-resistant *Staphylococcus aureus* and its emerging virulence. *Clin. Med. Res.* **3**, 57–60.
- Sidhu, M. S., Heir, E., Leegaard, T., Wiger, K., and Holck, A. (2002). Frequency of disinfectant resistance genes and genetic linkage with  $\beta$ -lactamase transposon Tn552 among clinical staphylococci. *Antimicrob. Agents Chemother.* **46**, 2797–2803.
- Sidhu, M. S., Heir, E., Sorum, H., and Holck, A. (2001). Genetic linkage between resistance to quaternary ammonium compounds and beta-lactam antibiotics in food-related *Staphylococcus* spp. *Microb. Drug Res. (Larchmont, N.Y.)* **7**, 363–371.
- Singer, R. S., Finch, R., Wegener, H. C., Bywater, R., Walters, J., and Lipsitch, M. (2003). Antibiotic resistance—The interplay between antibiotic use in animals and human beings. *Lancet Infect. Dis.* **3**, 47–51.
- Smalla, K., Heuer, H., Gotz, A., Niemeyer, D., Krogerrecklenfort, E., and Tietze, E. (2000). Exogenous isolation of antibiotic resistance plasmids from piggery manure slurries reveals a high prevalence and diversity of incq-like plasmids. *Appl. Environ. Microbiol.* **66**, 4854–4862.
- Smit, E., Wolters, A., and van Elsas, J. D. (1998). Self-transmissible mercury resistance plasmids with gene-mobilizing capacity in soil bacterial populations: influence of wheat roots and mercury addition. *Appl. Environ. Microbiol.* **64**, 1210–1219.
- Smith, S. R., Lang, N. L., Cheung, K. H., and Spanoudaki, K. (2005). Factors controlling pathogen destruction during anaerobic digestion of biowastes. *Waste Manag.* **25**, 417–425.
- Sorum, H., L'Abée-Lund, T. M., Solberg, A., and Wold, A. (2003). Integron-containing INCU R plasmids pRAS1 and pAr-32 from the fish pathogen *Aeromonas salmonicida*. *Antimicrob. Agents Chemother.* **47**, 1285–1290.
- Stokes, H. W., Holmes, A. J., Nield, B. S., Holley, M. P., Nevalainen, K. M., Mabbutt, B. C., and Gillings, M. R. (2001). Gene cassette PCR: Sequence-independent recovery of entire genes from environmental DNA. *Appl. Environ. Microbiol.* **67**, 5240–5246.
- Tennstedt, T., Szczepanowski, R., Krahn, I., Puhler, A., and Schluter, A. (2005). Sequence of the 68,869 bp incp-1alpha plasmid ptb11 from a waste-water treatment plant reveals a

- highly conserved backbone, a Tn402-like integron and other transposable elements. *Plasmid* **53**, 218–238.
- Tiemersma, E. W., Bronzwaer, S., Lyytikäinen, O., Degener, J. E., Schrijnemakers, P., Bruinisma, N., Monen, J., Witte, W., Grundmann, H., and Participants, U. K. E. A. R. S. S. (2004). Methicillin-resistant *Staphylococcus aureus* in Europe, 1999–2002. *Emerg. Infect. Dis.* **10**, 1627–1634.
- Tikofsky, L. L., Barlow, J. W., Santisteban, C., and Schukken, Y. H. (2003). A comparison of antimicrobial susceptibility patterns for *Staphylococcus aureus* in organic and conventional dairy herds. *Microb. Drug Res. (Larchmont, N.Y.)* **9**(Suppl 1), S39–S45.
- Tomlin, J., Pead, M. J., Lloyd, D. H., Howell, S., Hartmann, F., Jackson, H. A., and Muir, P. (1999). Methicillin-resistant *Staphylococcus aureus* infections in 11 dogs. *Vet. Rec.* **144**, 60–64.
- Top, E., De Smet, I., Verstraete, W., Dijkmans, R., and Mergeay, M. (1994). Exogenous isolation of mobilizing plasmids from polluted soils and sludges. *Appl. Environ. Microbiol.* **60**, 831–839.
- Turtura, G. C., Massa, S., and Ghazvinizadeh, H. (1990). Antibiotic resistance among coliform bacteria isolated from carcasses of commercially slaughtered chickens. *Int. J. Food Microbiol.* **11**, 351–354.
- Turutoglu, H., Ercelik, S., and Ozturk, D. (2006). Antibiotic resistance of *Staphylococcus aureus* and Coagulase-negative Staphylococci isolated from bovine mastitis. *Bull. Vet. Inst. Pulawy* **50**, 41–45.
- van Den Bogaard, A. E., London, N., and Stobberingh, E. E. (2000). Antimicrobial resistance in pig faecal samples from the Netherlands (five abattoirs) and Sweden. *J. Antimicrob. Chemother.* **45**, 663–671.
- van den Bogaard, A. E., and Stobberingh, E. E. (1999). Antibiotic usage in animals: impact on bacterial resistance and public health. *Drugs* **58**, 589–607.
- van Duijkeren, E., Box, A. T., Mulder, J., Wannet, W. J., Fluit, A. C., and Houwers, D. J. (2003). Methicillin resistant *Staphylococcus aureus* (MRSA) infection in a dog in the Netherlands. *Tijdschrift voor diergeneeskunde* **128**, 314–315.
- van Duijkeren, E., Wolfhagen, M. J., Box, A. T., Heck, M. E., Wannet, W. J., and Fluit, A. C. (2004). Human-to-dog transmission of methicillin-resistant *Staphylococcus aureus*. *Emerg. Infect. Dis.* **10**, 2235–2237.
- Vengust, M., Anderson, M. E., Rousseau, J., and Weese, J. S. (2006). Methicillin-resistant staphylococcal colonization in clinically normal dogs and horses in the community. *Letts. Appl. Microbiol.* **43**, 602–606.
- Voss, A., Loeffen, F., Bakker, J., Klaassen, C., and Wulf, M. (2005). Methicillin-resistant *Staphylococcus aureus* in pig farming. *Emerg. Infect. Dis.* **11**, 1965–1966.
- Waldron, D. E., and Lindsay, J. A. (2006). Sau1: A novel lineage-specific type I restriction-modification system that blocks horizontal gene transfer into *Staphylococcus aureus* and between *S. Aureus* isolates of different lineages. *J. Bacteriol.* **188**, 5578–5585.
- Weese, J. S., Archambault, M., Willey, B. M., Dick, H., Hearn, P., Kreiswirth, B. N., Said-Salim, B., McGeer, A., Likhoshvay, Y., and Prescott, J. F. (2005a). Methicillin-resistant *Staphylococcus aureus* in horses and horse personnel, 2000–2002. *Emerg. Infect. Dis.* **11**, 430–435.
- Weese, J. S., Rousseau, J., Traub-Dargatz, J. L., Willey, B. M., McGeer, A. J., and Low, D. E. (2005b). Community-associated methicillin-resistant *Staphylococcus aureus* in horses and humans who work with horses. *J. Am. Vet. Med. Assoc.* **226**, 580–583.
- Weese, J. S., Rousseau, J., Willey, B. M., Archambault, M., McGeer, A., and Low, D. E. (2006). Methicillin-resistant *Staphylococcus aureus* in horses at a veterinary teaching hospital: Frequency, characterization, and association with clinical disease. *J. Vet. Intern. Med.* **20**, 182–186.

- Weigel, L. M., Clewell, D. B., Gill, S. R., Clark, N. C., McDougal, L. K., Flannagan, S. E., Kolonay, J. F., Shetty, J., Killgore, G. E., and Tenover, F. C. (2003). Genetic analysis of a high-level vancomycin-resistant isolate of *Staphylococcus aureus*. *Science* **302**, 1569–1571.
- White, P. A., McIver, C. J., Deng, Y., and Rawlinson, W. D. (2000). Characterisation of two new gene cassettes, *aadA5* and *dfrA17*. *FEMS Microbiol. Lett.* **182**, 265–269.
- Witte, W. (1997). [How great is the potential danger from vancomycin-resistant enterococci?]. *Dtsch. Med. Wochenschr.* **122**, 1161–1163.
- Witte, W. (2000). Ecological impact of antibiotic use in animals on different complex microflora: environment. *Int. J. Antimicrob. Agents* **14**, 321–325.

CORRECTED PROOF